

SCIENCE BULLETIN OF JOSAI UNIVERSITY

城西大学理学部研究報告

Vol. 10 March 2002

FACULTY OF SCIENCE
JOSAI UNIVERSITY
SAKADO, SAITAMA, JAPAN

SCIENCE BULLETIN OF JOSAI UNIVERSITY

城西大学理学部研究報告

Vol. 10 March 2002

Editorial Board

Hikomichi Uehara (Chairman)

Hiroyuki Ishibashi

Norio Okada

Hiroshi Miyamae

Masakazu Akashi

Hideo Kagami

CONTENTS

PART I ANNUAL REPORTS

1. Abstracts of Papers Published in Journals	3
Physics	3
Chemistry	4
Physical Education	10
2. Books, Reviews and Other Printings	13
Mathematics	13
Chemistry	13
Biology	13
Earth Science	13
Physical Education	14
3. Oral Presentations	15
Mathematics	15
Chemistry	15
Physical Education	20



PART I ANNUAL REPORTS

1. Abstracts of Papers Published in Journals

PHYSICS

Absolute Photoionization Cross section Measurements of Xe^+ Ions in the 4d Threshold Energy Region

Yoh Itoh, Akira Ito^{*1}, Masashi Kitajima^{*2}, Tetsuo Koizumi^{*1}, Takao M. Kojima^{*3}, Hiroshi Sakai^{*1}, Mutsumi Sano^{*4} and Naoki Watanabe^{*5} (*1 Department of Physics, Rikkyo University, *2 Department of Physics, Sophia University, *3 The Institute of Physical and Chemical Research, *4 Japan Synchrotron Radiation Research Institute, *5 Institute of Low Temperature Science, Hokkaido University)

J. Phys. B: At. Mol. Opt. Phys., **34**, 3493–3499 (2001)

Using the photon-ion merged-beam technique, we have measured the absolute photoionization cross sections for Xe^{2+} and Xe^{3+} formation from Xe^+ at selected energies between 80 and 140 eV, and utilized the absolute data to normalize our previously reported relative cross sections. The structure and energy dependence of the cross sections for Xe^+ deviate from those of neutral Xe; however, the maximum total cross section for Xe^+ is, within the experimental error, the same as for Xe.

Relative Counting Efficiencies of Ion Charge-States by Microchannel Plate

S. Yagi^{*1} T. Nagata^{*1}, M. Koide^{*2}, Y. Itoh, T. Koizumi^{*3} and Y. Azuma^{*4} (*1 Department of Physics, Meisei University, *2 Institute for Laser Science, University of Electro-communications, *3 Department of Physics, Rikkyo University, *4 Photon Factory, KEK)

Nucl. Instrum. Methods Phys. Res. **B183**, 476–486 (2001)

Relative detection efficiencies D_{q+} of ions with different charges and impact energies in single-ion counting mode with microchannel plates (MCP, Hamamatsu F1094-21S) have been examined for Li^{q+} , Ar^{q+} , Ba^{p+} and Yb^{p+} ions ($q=1, 2$ and $p=1, 2, 3$). The incident ion energy ranges from 1.1 to 15 keV. The efficiencies D_{q+} for a given atom, plotted as a function of incident ion energy E_K lie on a unique curve. This comes originally from the fact that the secondary electron emission at the input channel-wall of MCP is dominated mainly by kinetic emission processes and little by potential emission processes in the keV energy range. When ions with different charges but the same E_K value of a given atomic species are detected, almost equal detection efficiencies are ensured as far as E_K is larger than 4 keV. Nearly equal detection efficiencies for singly and doubly charged ions accelerated by the same voltage are attainable at higher acceleration voltages for heavier ions, e.g., 5 kV for Ar^{2+} ions and about 10 kV for Ba^+ and Ba^{2+} ions.

CHEMISTRY

Evaluation of Expansion Coefficients from Optimal Fitting Parameters for the Analysis of Spectra of Diatomic Molecules and an Application to LiH

Hiromichi Uehara and J. F. Ogilvie* (*Centre for Experimental and Constructive Mathematics, Simon Fraser University)

J. Mol. Spectrosc., **207**, 143–152 (2001)

To evaluate individual expansion coefficients composing fitting parameters of the Born-Oppenheimer corrections to Dunham's coefficients Y_j that have been given analytically with the Δ_B and Δ_ω formalism, we examined the consistency of analytic expressions for those corrections with Watson's assertion of the experimental inseparability of non-adiabatic corrections $Q_{a,b}(r)$ for a molecule AB . Derived analytic expressions in terms of optimal fitting parameters for the corrections are essential to evaluate individual expansion coefficients. These expressions also reveal redundancies between empirical correction parameters Δ_{ij} . A method of evaluating nonadiabatic vibrational corrections $Q_{a,b}(r)$ and adiabatic corrections $S_{a,b}(r)$ separately consistent with Watson's assertion of inseparability is presented and is applied to an analysis of spectral data of LiH. Functions $Q_{a,b}$ and $S_{a,b}$ for LiH are thus successfully evaluated; $S_{H,Li}(r)$ values agree well with those predicted simply by wobble-stretch theory. Experimental values for optimal fitting parameters r_{1q}^H and r_{2q}^H are nearly equal to those of r_{1q}^{Li} and r_{2q}^{Li} , respectively, in agreement with a theoretical relation $r_{iq}^a = r_{iq}^b$.

Infrared Absorption Spectroscopy of $Ar-^{12}C^{18}O_2$: Change in the Intramolecular Potential upon Complex Formation

Yasushi Ozaki, Kouji Horiai, Toichi Konno, and Hiromichi Uehara

Chem. Phys. Lett., **335**, 188–194 (2001)

An infrared absorption spectrum has been observed and analyzed for an $Ar-^{12}C^{18}O_2$ van der Waals (vdW) complex in the range of 2312–2314 cm^{-1} . The effective structure of the complex derived from the obtained rotational constants is in essential agreement with the structure estimated from that of $Ar-^{12}C^{16}O_2$ based on the isotope effect on the amplitudes of vdW vibrations. The band origin for the $Ar-^{12}C^{18}O_2$ ν_4 mode shows a red shift of 0.4567 cm^{-1} with respect to that for $^{12}C^{18}O_2$ ν_3 mode; the change in the intramolecular potential of CO_2 upon complex formation with Ar is determined from the isotope effect on the band-origin shift.

Comparative Assessment of the Efficiency of Thin Film Electrodes Fabricated by Three Deposition Methods

Photoelectrochemical Degradation of the DBS Anionic Surfactant

Hisao Hidaka*¹, Kazuhiko Ajisaka*¹, Satoshi Horikoshi*¹, Toshiyuki Oyama, Kazuo Takeuchi*², Jincal Zhao*³, Nick Serpone*⁴ (*1 Frontier Research Center for the Global Environmental Protection, Meisei University, *2 The Institute of Physical and Chemical Research (Riken), *3 Institute of Chemistry, Chinese Academy of Sciences, *4 Department of Chemistry and Biochemistry, Concordia University)

J. Photochem. Photobiol. A: Chem. **138**, 185–192(2001)

This study reports on a comparative assessment of the efficiency of TiO₂/SnO₂-doped thin film electrodes fabricated by three different deposition methods. Process efficiencies, as measured by the kinetics of degradation, were examined by monitoring the photooxidative fate of two test substrates, namely the anionic surfactant sodium dodecylbenzenesulfonate (DBS) and the model compound sodium benzenesulfonate (BS). Methods chosen to fabricate the TiO₂/OTE electrodes comprised (1) pulsed laser deposition, (2) deposition of a TiO₂ paste on the OTE plate, and (3) sol-gel deposition. Both DBS and BS are readily photo-oxidized and, under our conditions, were anodically biased at +0.3 V, which generated a photocurrent during the photoelectrolysis. The sol-gel (C) and anatase (B1) electrodes proved especially useful under the +0.3 V bias. In the absence of any bias, however, DBS photodegraded faster on the TiO₂ pasted electrode. Under otherwise similar conditions, pulsed laser rutile TiO₂/OTE electrode showed very little photoactivity. Some of the characteristics of the TiO₂ film, such as the TiO₂ crystalline form, film thickness, surface roughness, and film transparency appear to have some effect on the overall photodegradative process efficiency.

Photooxidative *N*-demethylation of Methylene Blue in Aqueous TiO₂ Dispersions under UV-Irradiation

Tianyong Zhang*¹, Toshiyuki Oyama, Akeo Aoshima*¹, Hisao Hidaka*¹, Jincal*², and Nick Serpone*³ (*1 Frontier Research Center for the Global Environmental Protection, Meisei University, *2 Institute of Chemistry, Chinese Academy of Sciences, *3 Department of Chemistry and Biochemistry, Concordia University)

J. Photochem. Photobiol. A: Chem. **140**, 163–172 (2001)

Methylene blue (MB) is a representative of a class of dyestuffs resistant to biodegradation. Its decomposition was examined in aqueous TiO₂ dispersions under UV-illumination to assess the influence of temperature, pH, concentration of dissolved oxygen (DOC), initial concentration of MB, and light intensity on the kinetics of decomposition. Hypsochromic effects (i.e. blue shifts of spectral bands) resulting from *N*-demethylation of the dimethylamino group in MB occurs concomitantly with oxidative degradation. Measured at constant pH 4 were also the maximum quantity of MB adsorbed on TiO₂, and the

kinetics of degradation of MB and of TOC removal. Photobleaching of MB solutions takes place at low DOCs and is caused by a reversible reductive process involving photogenerated electrons on TiO_2 . The rate of degradation of MB remains fairly constant regardless of whether the dispersion was purged with oxygen prior to irradiation or with air during the light irradiation period. The photocatalytic process depends on light intensity, but not on the total light energy absorbed. The photoreaction followed pseudo first-order kinetics even at high MB concentrations (0.3 mM). The temperature dependence of the photodegradation kinetics was assessed ($E_a=8.9$ kJ/mol), as well as the relative photonic efficiency, ξ_r , relative to phenol (0.4).

Assessment and Influence of Operational Parameters on the TiO_2 Photocatalytic Degradation of Sodium Benzene Sulfonate Under Highly Concentrated Solar Light Illumination

Tianyong Zhang*¹, Toshiyuki Oyama, Satoshi Horikoshi*¹, Jincui Zhao*², Hisao Hidaka*¹, and Nick Serpone*³, (*1 Frontier Research Center for the Global Environmental Protection, Meisei University, *2 Institute of Chemistry, Chinese Academy of Sciences, *3 Department of Chemistry and Biochemistry, Concordia University)

Solar Energy, **71**, 305–313 (2001)

Sodium benzene sulfonate (BS) was decomposed in aqueous TiO_2 dispersions under highly concentrated solar light illumination to examine the photocatalytic characteristics of a parabolic round concentrator (PRO) reactor to degrade the pollutant without visible light absorption. The effects of such operational parameters as initial concentration, volume of the aqueous BS solution, oxygen purging, and TiO_2 loading on the kinetics of decomposition of BS were investigated. An effective photodegradation necessitates a suitable combination of initial volume and concentration of BS solution. Relative to atmospheric air, oxygen purging significantly accelerates the degradation process at high initial concentrations of BS (0.40 mM or 1.0 mM). Optimal TiO_2 loading was 9 g litre⁻¹, greater than previously reported. Elimination of TOC (total organic carbon) followed pseudo first-order kinetics in the initial stages of the photodegradation process. The relative photonic efficiency for the photodegradation of BS is $\zeta_{\text{rel}} = 1.0$.

Syntheses of Tetraazabis(tropocoronand)s Containing Hydroxy Group(s) and Conformational Analysis of Their Nickel(II) Complexes in Solution

Ohki Sato*¹, Hideki Chikamatsu*¹, Josuke Tsunetsugu*¹, Kimio Shindo, Hidetsugu Wakabayashi, and (the late) Tetsuo Nozoe*² (*1 Department of Chemistry, Faculty of Science, Saitama University, *2 Tokyo Research Laboratories, Kao Corporation)

Heterocycles, **52** (1), 459–463 (2000)

The stepwise syntheses of one symmetric and three asymmetric tetraazabis-

(tropocoronand)s (**12a–12d**) having two linker chains, with or without a hydroxy group, of the same or different length are reported. The coordination styles of their nickel(II) complexes (**13a–13d**) in solution have also been discussed based on their ^1H NMR and UV-VIS. spectral data.

Synthesis and Properties of Tetraazazulenocoronands and Octaazazulenocryptands

Hidetsugu Wakabayashi, Teruo Kurihara, and (the late) Tetsuo Nozoe* (*Tokyo Research Laboratories, Kao Corporation)

Heterocycles, **54** (1), 87–92 (2001)

Treatment of 1, 3-diformylazulene (**1**) with 1.2 equiv. of α, ω -alkanediamines (**3** and **4**) and tris (2-aminoethyl) amine (**5a**) in ethanol for 24 h at room temperature gave tetraazaazulenecoronands (**2** and **9**) and octaazaazulenocryptand (**10**) in one-pot procedure in good yields.

Convenient Synthesis of 4-Trifluoromethyl-Substituted Imidazole Derivatives

Masami Kawase*¹, Setsuo Saito*¹, and Teruo Kurihara (*¹ Faculty of Pharmaceutical Sciences, Josai University)

Chem. Pharm. Bull., **49** (4), 461 (2001)

Mesoionic 4-trifluoroacetyl-1, 3-oxazolium-5-olates (**1**), obtained from the reaction of *N*-acyl-*N*-alkylglycines (**2**) with trifluoroacetic anhydride, react with ammonia to give 4-trifluoromethyl-3, 4-dihydroimidazoles (**3**) in high yields. Dehydration of **3** gives 4-trifluoromethylimidazoles (**4**) in high yields. The novel ring transformation of **1** into **3** occurs via a regioselective attack of ammonia on the C-2 position of the ring.

Biological Activity of a Fruit Vegetable, “Anastasia Green”, a Species of Sweet Pepper

Noboru Motohashi*¹, Teruo Kurihara, Hidetsugu Wakabayashi, Mikako Yaji, Ilona Mucsi*², Joseph Molnar*², Shichiro Maruyama*³, Hiroshi Sakagami*³, Hideki Nakashima*⁴, Satoru Tani*⁵, Yoshiaki Shirataki*⁵, and Masami Kawase*⁵ (*¹ Meiji Pharmaceutical University, *² Faculty of Medicine, Institute of Microbiology, Albert Szent-Gyorgyi Medical University, *³ Department of Dental Pharmacology, Meikai University School of Dentistry, *⁴ Department of Microbiology and Immunology, Kagoshima University Dental School, *⁵ Faculty of Pharmaceutical Sciences, Josai University)

In vivo., **15**, 437–442 (2001)

Russian green sweet pepper (Anastasia Green) was successively extracted with hexane, acetone, methanol and 70% methanol and the extracts were further separated into a total of twenty fractions by silica gel or ODS column chromatographies. The extracts and fractions showed higher cytotoxic activity against two human oral tumor cell lines than

against normal human gingival fibroblasts, suggesting their tumor-specific action. Several fractions [H3, H4, A4] reversed the multidrug resistant gene (MDRI) against L5178 mouse T-cell lymphoma more effectively than (\pm) verapamil (positive control). All extracts and fractions showed no anti-human immunodeficiency virus (HIV) nor anti-*Helicobacter pylori* activity. These data suggest the medicinal importance of an Anastasia Green extract.

Biological Activity of kiwifruit Peel Extracts

Noboru Motohashi*¹, Yoshiaki Shirataki*², and Masami Kawase*², Satoru Tani*², Hiroshi Sakagami*³, Kazue Satoh*⁴, Teruo Kurihara, Hideki Nakashima*⁵, Kristina Wolfard*⁶, Csilla Miskolci*⁶, and Joseph Molner*⁶ (*1 Meiji Pharmaceutical University, *2 Faculty of Pharmaceutical Sciences, Josai University, *3 Department of Dental Pharmacology, Meikai University School of Dentistry, *4 Analysis Center, School of Pharmaceutical Sciences, Showa University, *5 Department of Microbiology and Immunology, Kagoshima University Dental School, *6 Faculty of Medicine, Institute of Microbiology, Albert Szent-Gyorgyi Medical Univeresity)

Phytother. Res., **15**, 337–343 (2001)

Various bioactive substances in Kiwifruit extracts were fractionated by organic solvent extractions, followed by silica gel and ODS chromatographies. Both cytotoxic activity and multi-drug resistance reversal activity were found in the less polar fractions. Cytotoxic activity was not always parallel the radical intensity. Antibacterial activity was distributed into various fractions and all fractions were inactive against *Candida albicans* and *H. pylori*. Only 70% methanol extracts showed anti-human immunodeficiency virus activity, and produced a broad ESR signal under alkaline conditions, in a fashion similar to lignin. These fractions also effectively scavenged O₂⁻ produced by the xanthine-xanthine oxidase reaction, suggesting a bimodal (pro-oxidant and antioxidant) action. These data suggest a medicinal efficacy of kiwifruit peel extracts.

Calorimetric study of nickel complex compounds with tmen as a ligand: determination of the energetic parameters for the δ - λ conformational change of the ethylenediamine framework

M. Oguni*¹, T. M. Yoshida*¹, K. Wada*¹, Y. Fukuda*², N. Ogino, T. Ito, H. Miyamae, K. Sato*³ (*1 Department of Chemistry, Faculty of Science, Tokyo Institute of Technology, *2 Department of Chemistry, Faculty of Science, Ochanomizu University, *3 Department of Chemistry, Faculty of Science, Niigata University)

J. Phys. Chem. Solids, **62**, 613–618 (2001)

Heat capacities of crystalline $[\text{Ni}(\text{acac})(\text{tmen})(\text{H}_2\text{O})_2]\text{I}$ and $[(\text{Ni}(\text{acac})_2(\text{tmen}))]$ were measured with an adiabatic calorimeter in the range 10–300 K, where acac and tmen are acetylacetonate and *N, N, N', N'*-tetramethylenediamine, respectively. Glass transitions

were found at (108 ± 1) and (126 ± 1) K for the respective compounds. They were interpreted as due to the freezing-in of disorder between the δ and λ conformations of tmen moiety. The energy difference ($\Delta\epsilon$) between the two conformations and the activation energy ($\Delta\epsilon_a$) for the conformational change were determined by the analyses of the calorimetric data to be (0.87 ± 0.05) and (32 ± 1) kJ mol⁻¹ for the iodide compound, and (1.58 ± 0.05) and (39 ± 1) kJ mol⁻¹ for the molecular compound, respectively. It is discussed that $\Delta\epsilon$ is affected both by the coordination structure of complex and by the presence/absence and species of counter anion while that $\Delta\epsilon_a$ is determined rather by the interaction of tmen with the other ligands within the complex. X-ray diffraction technique was applied to crystalline $[\text{Ni}(\text{acac})(\text{tmen})(\text{H}_2\text{O})_2] \text{ClO}_4$ to confirm the above interpretation, and the details are given as an appendix.

Heterocyclization of 4-trifluoroacetyl-1, 3-oxazolium-5-olates with 1, 4-bis-nucleophiles

Masami Kawase*¹, Hiromi Koiwai*¹, Toru Tanaka*¹, Satoru Tani*¹, and Hiroshi Miyamae (*1 Faculty of Pharmaceutical Sciences, Josai University)

Heterocycles, **55**, 1919–1926 (2001)

Reactions of aromatic 1,4-bis-nucleophiles such as *o*-phenylenediamine and *o*-aminothiophenol, with mesoionic 4-trifluoroacetyl-1,3-oxazolium-5-olates(**1**) gave regiospecifically seven member trifluoromethylated heterocycles such as 1,5-benzodiazepines(**3**) and 1,5-benzothiazepines(**4**). The reaction with *o*-aminophenol afforded non-cyclized products(**5**). The structures of **3**, **4**, and **5** were established by X-Ray diffraction analysis.

Changes in Amino Acid Pool and Utilization during Apoptosis in HL60 Cells Induced by Epigallocatechin Gallate or Gallic Acid

Hiroshi Sakagami*¹, Yoshiko Yokote, Kiso Akahane (*1 Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama)

Anticancer Res., **20**, 265–270 (2001)

Recent studies have demonstrated the apoptosis-inducing potential of epigallocatechin gallate (EGCG), a major component of green tea, against various cultured cell lines. By using an amino acid analyzer, we investigated here the possible changes in the amino acid pool and utilization during the apoptosis of HL-60 cells induced by EGCG or gallic acid, a structural unit of tannin. Sublethal concentrations of EGCG initially elevated and then reduced the intracellular concentrations of most of amino acids except for glutamic acid and aspartic acid, whereas lethal concentrations of EGCG continuously reduced these amino acid pools during 6 hours. Both sublethal and lethal concentrations of gallic acid initially elevated and then reduced these amino acid pools. Both inducers elevated the intracellular accumulation and production of arginine and extensively reduced the utilization of other amino acids. These data demonstrate that EGCG showed more severe effects on the amino

acid pool and utilization than gallic acid, which may explain, at least in part, the difference in apoptosis-inducing potential between these inducers.

PHYSICAL EDUCATION

A Study of Line-up Analysis in International Women's Volleyball Game's —2000 Olympic Final Qualifications: Japanese Team versus Croatian Team—

Daisen Shimazu*¹, Kyoichi Izumikawa*², Sotonori Yamamoto*³, Masakazu Akashi*⁴, Mituru Sakai*⁵, Takehiko Tahara*⁶, Satoshi Harada*⁷ (*1 Japan Women's University, Department of Physical Education, *2 Kanagawa Institute of Technology, Department of Physical Education, *3 Kyorin University, Department of Physical Education, *4 Josai University, Department of Physical Education, *5 Kyushu Women's Junior College, Department of Physical Education, *6 Nara University, Department of Physical Education, *7 Rishou University, Department of Physical Education)

J. Phy. Ex. Sports., 7, 11-20 (2001)

For the Japanese, the probable higher-rank groups were Line-up5 (L-up 5) and 1, while the line-ups to be avoided were L-up2 and 4. For L-up5, which took first place in the estimated total ranking, the best starting rotation phase was expected to be rotation Phase3 (R3) if the Japanese team was on the serving side or R2 if on the receiving side, provided that the Croatian team starts with its rotation R6. With this line-up, it was believed to be essential for the Japanese team to enhance the defensive power of R5, which would give a lower defensive power index, and to maintain the offensive power of R1, which it would give higher offensive power index. It was also required that the team improve the offensive power of R5, which have lower offensive power index when compared to R1, as well as to retain defensive power for R2 and R3 with higher defensive power indexes.

The Japanese team selected L-up 1 and 6 instead of Line-up5 for the match. Line-up1 usually decreases in defensive power index for R3 and R4. These rotation phases chosen by Japan faced R3 and R4 of the Croatian L-up2, as well as, Line-up6 usually decreases in defensive power index for R4. This rotation phases chosen by Japan faced R3 of the Croatian L-up3, which were estimated to be higher in offensive power index. The Japanese team failed to enhance the inferior defensive power of L-up 1 and 6 for these rotation phases, while the Croatian team successfully retained and even improved the strong offensive power for the corresponding rotation phases. We judged that this fact was one of the reasons why the Japanese team failed to win the match. Therefore the next challenge to be addressed should be an analysis of line-ups for cases in which the opposing team is on the servicing side.

This was a guide of volleyball game analysis by strong or weak power of rotation phase and contributive rate of individual player's points or side-out points. But it has need to develop.

Key words; Volleyball, Game analysis, Offensive Power index. Defensive Power index,

B-T model

A Study of Line-up Analysis in International Women's Volleyball Games — '99 World Cup Women's Games, Japan Team versus China Team —

Daisen Shimazu*¹, Kyoichi Izumikawa*², Sotonori Yamamoto*³, Masakazu Akasi*⁴, Mituru Sakai*⁵, Takehiko Tahara*⁶, Satoshi Harada*⁷ (*1 Japan Women's University, Department of Physical Education, *2 Kanagawa Institute of Technology, Department of Physical Education, *3 Kyorin University, Department of Physical Education, *4 Josai University, Department of Physical Education, *5 Kyushu Women's Junior College, Department of Physical Education, *6 Nara University, Department of Physical Education, *7 Rishou University, Department of Physical Education)

Journal of Volleyball Science, 3-1, 15-25 (2001)

The authors attempted to extrapolate the best and worst line-up for the Japan women's volleyball team in the '00 Olympic Final Qualification games with China. They used the maximum likelihood method and Brodley-Terry Model to retrospectively investigate the nine games between Japan, Cuba, Russia, Brazil, and China in the '99 World Cup Women's Volleyball Games held in Japan (from November 2 to 16, 66 games in total).

Line-up4 was obtained by the best and Line-up6 was the worst. The starting line-up of the Japan team in line-up4 was R5 with and without the serve. In Line-up6, the starting line-up was R3 with and without the serve, respectively. Line-up6 was the line-up that the Japan team should have avoided. To enhance the team's performance, it was necessary to improve the rotation phase with a low offense index and defense index. High defense index was necessary to maintain. Specifically, R4 R5 and R6 (low offense index), R1 and R2 (low defense index), R3, R5 and R6 (high defense index) were applicable to rotation phases for the Japan team in Line-up4. R6 (18.5%) and R2 (-13) were obtained the most low offense performance by point rate and Net Score each method.

Key words; Volleyball, Game analysis, Offense index, Defense index

Effects of intake of foods with antioxidant activities in long distance runners.

Kojiro Ishii*¹, Kazuki Takizawa*¹, Kazuya Yonezawa*², Eiji Daigo*³, Noto Suzuki*⁴ (*1 Grad. School of Edu., Hokkaido Univ., *2 Grad School of Med, hokkaido Univ., *3 Kanagawa Univ., *4 Josai Univ.) & Teruaki Komine (Noevlr Foods Co., Ltd.)

Advances in Exercise and Sports Physiology, 6 (4) 160 (2000)

<Objective>

As the oxygen consumption increases due to exercise. The production of active oxygen increases in the body. Hyperoxidation has been reported to cause skeletal muscle injuries, hemolysis, and DNA damages, and intake of foods with antioxidant activities is recommended. In this study oxidation stress in long distance runners during the training

period including a training camp was examined, and the effects of intake of functional food items (fermented grain extracts) with antioxidant activities (AOZ) were evaluated.

〈Methods〉

Study 1: The cholesteryl ester hydroperoxide (CEOOH) and 8-hydroxy-deoxyguanosine (8-OHdG) levels were measured in 14 female college long-distance runners, who were divided into early intake and late intake groups.

Study 2: The CPK, RBC, Ht, Hb, and Fe levels were measured in 29 male long distance runners, who were divided into AOZ and placebo groups.

〈Results and Discussion〉

The CEOOH level was increased at the end of the training camp in both the early intake and late intake groups, indicating accumulation of oxidation stress during the training camp.

Ht, Hb, and Fe showed interaction (intake or no intake X intake period), and the blood profile remained stable in the AOZ group even during the training camp.

Active oxygen produced during intensive endurance training is known to surpass the ability of its elimination. Intake of foods with antioxidant activities is considered to be a preventive measure against oxidation stress.

〈Keyword〉

Endurance Running, Training, Reactive Oxygen Species (ROS)

2. Books, Reviews and Other Printing

MATHEMATICS

Invariant hyperfunctions on the prehomogeneous vector space acting the group $GL_n(\mathbf{R}) \times SO_{p,q}(\mathbf{R})$.

Representation theory and noncommutative harmonic analysis (Japanese) (Kyoto, 1998)
M. Muro: Sūrikaiseikikenkyūsho Kōkyūroku, No. 1082, 93-101 (1999) by T. Kogiso in *Mathematical Reviews of American Math. Soc.* (2000)

CHEMISTRY

Separation of DNA by Free Flow Electrophoresis in Space

Hidesaburo Kobayashi and Naoaki Ishii
Biological Sciences in Space Vol. 15, Supplement, s195. (2001)

BIOLOGY

キイロシヨウジヨウバエの成虫体重に関する人為選抜及びその生存力に与える影響

小須田和彦
 城西大学研究年報 (自然科学編), 24, 1-10

EARTH SCIENCE

南極周辺の大陸縁辺域の形成過程

加賀美英雄, 森脇喜一*, 野木義史* (*国立極地研究所)
 平成 10 年度共同研究報告書
 国立極地研究所 pp. 74-75 (2001)

関東山地四万十帯のホルンヘルス IV-塩山市泉水谷の泥質岩変成鉱物の産状と化学組成

加賀美英雄, 谷口英嗣* (*駒沢大学高等学校)
 城西大学研究年報 (自然科学編) 25, 1-37 (2001)

アルプスの形成とその岩石

加賀美英雄

季刊水路 30, 2, 18-22 (2001)

PHYSICAL EDUCATION**バレーボール選手の体力に関する研究—城西大学選手について—**

明石正和, 古泉一久*, 千葉 正**, 塚田 勇*** (*城西大学非常勤講師, **修紅短期大学, ***城西医療技術専門学校理学療法学科)

城西大学研究年報・自然科学編, 25, 2001年12月, pp.53-61

エアロビクス・ダンス運動による生理学的変動の検討

中尾喜久子*, 横内靖典, 永都久典, 鈴木尚人, 古田裕子**, 林貢一郎***, 相澤勝治***, 目崎 登*** (*城西大学非常勤講師, **早稲田大学体育局非常勤講師, ***筑波大学)

城西大学研究年報・自然科学編, 25, 2001年12月, pp.63-70

全日本大学女子駅伝優勝チームの身体的特徴および体力について (第1報)

鈴木尚人, 上田 大* (*日本体育大学)

城西大学研究年報・自然科学編, 25, 2001年12月, pp.71-81

3. Oral Presentations

MATHEMATICS

解析的測度の絶対連続性について

山口 博

実解析学シンポジウム (東海大学), 2001 年 11 月

Convergence to Periodic Fronts in Nonlinear Diffusion Equations

Toshiko Ogiwara

第18回九州における偏微分方程式研究集会 (九州大学), 2001 年 1 月

Asymptotic Behavior of Solution for Nonlinear Diffusion Equations

Toshiko Ogiwara

Third Summer Workshop on Semilinear Elliptic Problems (Cheju, Korea), 2001 年 7 月

Asymptotic Behavior of Solution for Nonlinear Diffusion Equations

Toshiko Ogiwara

International Conference on Nonlinear Analysis and Convex Analysis (Hirosaki, Japan),
2001 年 7 月

Asymptotic Behavior of Solutions to a Model of Spiral Crystal Growth

Toshiko Ogiwara and Ken-Ichi Nakamura* (*Department of Computer Science, University
of Electro-Communications)

研究集会「非線形拡散系とその周辺」(京都大学数理解析研究所), 2001 年 10 月

CHEMISTRY

二原子分子のスペクトル解析から決定できる分子パラメーターと物理量

上原博通

分子構造総合討論会 (札幌), 2001 年 9 月, 講演要旨集 p. 600

KHの振動回転スペクトルと分子パラメーターの決定

堀合公威, 並木恵一*, 上原博通 (*長岡技術科学大学)

分子構造総合討論会 (札幌), 2001年9月, 講演要旨集 p. 601

(N₂O)₂ および N₂O-CO₂ 赤外ダイオードレーザー分光—錯体形成による N₂O ν₃ 振動の振動数変化—

尾崎 裕, 堀合公威, 紺野東一, 上原博通

分子構造総合討論会 (札幌), 2001年9月, 講演要旨集 p. 602

Change in the Intramolecular Potential of CO₂ upon Complex Formation with Ar

Yasushi Ozaki, Kouji Horiai, Toichi Konno, and Hiromichi Uehara

Spectroscopy in the 21st Century (Hayama, Japan), March 18–22, 2001, Abstracts, p. 112

粘度の新しい高精度測定法

末岡一生

日本化学会第79春季年会 (神戸), 2001年3月, 講演要旨集 I, 2D639.

Comparative Assessment of Photocurrent Efficiency of TiO₂/OTE Thin Film Electrodes, Fabricated by Three Deposition Methods, in the Photodegradation of Water-soluble Organic Materials

Hisao Hidaka*¹, Toshiyuki Oyama, Satoshi Horikoshi*¹, Jincal Zhao*², Nick Serpone*³
(*1 Frontier Research Center for the Global Environmental Protection, Meisei University,
*2 Institute of Chemistry, Chinese Academy of Sciences, *3 Department of Chemistry and
Biochemistry, Concordia University)

First International Conference on Semiconductor Photochemistry University of Strathclyde,
Glasgow (U.K.), July 23–25, 2001, Abstracts, p. 64

集光照射装置を用いた太陽光によるベンゼンスルホン酸ナトリウムの光触媒分解

大山俊之, 張 天永*¹, 堀越 智*¹, 趙 進才*², N. セルフォン*³, 日高久夫*¹ (*1 明星大学
地球環境保全センター, *2 Institute of Chemistry, Chinese Academy of Sciences, *3 Department
of Chemistry and Biochemistry, Concordia University)

第54回コロイドおよび界面化学討論会 (日野), 2001年9月, 講演要旨集, p. 182

Photocatalyzed *N*-demethylation and Degradation of Methylene Blue in Titania Dispersions Exposed to Concentrated Sunlight.

Tianyong Zhang^{*1}, Toshiyuki Oyama, Satoshi Horikoshi^{*1}, Jincal Zhao^{*2}, Nick Serpone^{*3}, and Hisao Hidaka^{*1} (*1 Frontier Research Center for the Global Environmental Protection, Meisei University, *2 Institute of Chemistry, Chinese Academy of Sciences, *3 Department of Chemistry and Biochemistry, Concordia University)

第 54 回コロイドおよび界面化学討論会 (日野), 2001 年 9 月, 講演要旨集, p. 194

テトラアザアズレノコロナンドの合成と性状

若林英嗣, 新藤君男, 栗原照夫, (故)野副鉄男^{*1} (*1 花王東研)
日本化学会第 78 回春季年会 (日大理工学部), 2000 年 3 月, 3P153

テトラアザトロポクリプタンドの合成と性状

新藤君男, 三村康幸, 若林英嗣, 栗原照夫
日本化学会第 78 回春季年会 (日大理工学部), 2000 年 3 月, 3P154

Biological Activity of Fruite Vegetable, “Anastasia Green”, A Species of Sweet Pepper

N. Motohashi^{*1}, T. Kurihara, H. Wakabayashi, M. Yaji, K. Wolfard^{*2}, J. Molnar^{*2}, S. Maruyama^{*3}, H. Sakagami^{*3}, H. Nakashima^{*4}, S. Tani^{*5}, Y. Shirataki^{*5}, and M. Kawase^{*5} (*1 Meiji Pharmaceutical University, *2 Faculty of Medicine, Institute of Microbiology, Albert Szent-Gyorgyi Medical University, *3 Department of Dental Pharmacology, Meikai University School of Dentistry, *4 Department of Microbiology and Immunology, Kagoshima University Dental School, *5 Faculty of Pharmaceutical Sciences, Josai University)
New Anticancer Agents (Athens, Greece), June. 9-12, 2001.

Biological Activity of Fruite Vegetable, “Anastasia Red”, A Species of Sweet Pepper

N. Motohashi^{*1}, H. Wakabayashi, T. Kurihara, Y. Takada, S. Maruyama^{*2}, H. Sakagami^{*2}, H. Nakashima^{*3}, S. Tani^{*4}, Y. Shirataki^{*4}, M. Kawase^{*4}, K. Wolfard^{*5}, and J. Molnar^{*5} (*1 Meiji Pharmaceutical University, *2 Department of Dental Pharmacology, Meikai University School of Dentistry, *3 Department of Microbiology and Immunology, Kagoshima University Dental School, *4 Faculty of Pharmaceutical Sciences, Josai University, *5 Faculty of Medicine, Institute of Microbiology, Albert Szent-Gyorgyi Medical University)
New Anticancer Agents (Athens, Greece), June. 9-12, 2001.

ヘテロ環が縮環したアズレンキノン誘導体の合成

若林英嗣, 松丸聖一, 小笠原聖, 栗原照夫, 安並正文*¹ (*¹ 日大工)

第 31 回構造有機化学討論会 (山口大学平川キャンパス), 2001 年 10 月, 2p.05

水熱合成できた [Ni(323-tet)]PbBr₄ と混合溶媒から析出した [Ni(323-tet)]PbCl₄

宮前 博, 佐藤惣一, 日原五郎

日本化学会第 79 春季年会 (神戸), 2001 年, 講演要旨集 I, p. 258

トリス(ポリピリジン)ルテニウム(II)とアスコルビン酸イオンとの光反応

立屋敷 哲*¹, 諸岡樹興, 木幡浩昌*¹, 篠崎一英*², 宮前 博, 日原五郎 (*¹ 女子栄養大学, *² 横浜市立大学大学院理学研究科)

第 14 回配位化合物の光化学討論会, 札幌 (北大学術交流会館), 講演要旨集, 2001 年 7 月, pp. 67-68.

[M(en)₂]Br₂ (M = Ni or Cu) と臭化鉛から出来た積層構造の再検討—SPRING-8 を利用した微小結晶構造解析—

宮前 博, 児玉竜也, 日原五郎, 本間法敬*¹, 飯田昭文*¹, 石井彰人*¹, 尾関智二*¹ (*¹ 東京工業大学理学部)

第 51 回錯体化学討論会 (松江), 講演要旨集, 2001 年 9 月, p. 221

トリス(ポリピリジン)ルテニウム(II)錯体の還元消光に伴う消光剤と配位子との反応

立屋敷 哲*¹, 諸岡樹興, 木幡浩昌*¹, 篠崎一英*², 宮前 博, 日原五郎 (*¹ 女子栄養大学, *² 横浜市立大学大学院理学研究科)

第 51 回錯体化学討論会 (松江), 講演要旨集, 2001 年 9 月, p. 684

混合摩砕によるジメチルグリオキシムとニッケル(II)塩との錯体の無溶媒合成

日原五郎, 佐藤雅浩, 宮前 博

第 12 回固体の反応性討論会講演要旨集 (東京), 2001 年 11 月, p. 71

ハト羽毛ケラチン遺伝子群の塩基配列について

高橋理恵子, 新井邦男, 横手よし子, 赤羽 競

第 74 回日本生化学会大会 (横浜), 2001 年 10 月, 講演要旨集, p. 951

円二色性 (CD) によるタンパク質の二次構造解析—Spectral Profile の特徴により分類した Reference の作成—

新井邦男, 室田浩一, 横手よし子, 高橋理恵子, 赤羽 競
第 74 回日本生化学会大会 (横浜), 2001 年 10 月, 講演要旨集, p. 956

マイクロフリーフロー電気泳動モジュールを用いたゾーン電気泳動におけるタンパク分離挙動

目黒一輝, 小林英三郎, 太田和希, 長濱久花, 田島信芳*, 篠原悦夫* (* オリンパス光学工業(株))
第 8 回クロマトグラフィーシンポジウム (東京), 2001 年 6 月, CHROMATOGRAPHY, Vol. 22, pp. 9-10

マイクロフリーフロー電気泳動モジュールを用いたタンパク分離

小林英三郎, 太田和希, 目黒一輝, 長濱久花, 田島信芳, 篠原悦夫* (* オリンパス光学工業(株))
Separation Sciences 2001 (東京), 2000 年 6 月, 講演要旨集

分子生物学と電気泳動

小林英三郎
Separation Sciences 2001 (東京), 2000 年 6 月, 講演要旨集

モノリス/C18 クロマトカラムにおけるタンパク分離挙動

小林英三郎, 松崎正博, 村上裕一
第 12 回クロマトグラフィー科学会議 (京都), 1999 年 9 月, CHROMATOGRAPHY, Vol. 22, 要旨集, pp. 91-92

マイクロフリーフロー電気泳動モジュールにおける電気浸透流測定

小林英三郎, 島村克好, 田島信芳*, 船崎 純*, 篠原悦夫* (* オリンパス光学工業(株))
第 21 回キャピラリー電気泳動シンポジウム 2001 (神戸), 2001 年 12 月, 要旨集, pp. 43-44

マイクロフリーフロー電気泳動モジュールにおける色素化合物の実効移動度測定

島村克好, 小林英三郎, 田島信芳*, 船崎 純*, 篠原悦夫* (* オリンパス光学工業(株))
第 21 回キャピラリー電気泳動シンポジウム 2001 (神戸), 2001 年 12 月, 要旨集, pp. 103-104

マイクロフリーフロー電気泳動モジュールにおける色素タンパクの実効移動度測定

小林英三郎, 島村克好, 田島信芳*, 船崎 純*, 篠原悦夫* (* オリンパス光学工業(株))
第 21 回キャピラリー電気泳動シンポジウム 2001 (神戸), 2001 年 12 月, 要旨集, pp.105-106

クロシヨウジョウバエ Acph 遺伝子の解析

北川浩子

第 73 回日本遺伝学会 (東京), 2001 年 9 月, Gene. Genet. Sys. **76**, p. 464.

PHYSICAL EDUCATION**全日本大学女子駅伝優勝チームの身体的特徴および体力について**

上田 大*¹, 森田修一*¹, 石井友保*¹, 鈴木尚人, 池原忠明*¹, 黄 仁官*¹, 堀居 昭*¹
(*¹ 日本体育大学)

第 56 回日本体力医学会 (仙台), 2001 年 9 月, p. 981

国際女子バレーボール試合のラインアップ分析—'00 シドニーオリンピック最終予選, 日本対イタリアの分析—

島津大宣*¹, 泉川喬一*², 山本外憲*³, 田原武彦*⁴, 坂井 充*⁵, 明石正和, 原田 智*⁶ (*¹ 日本女子大学, *² 神奈川工科大学, *³ 杏林大学, *⁴ 奈良大学, *⁵ 九州女子短期大学, *⁶ 立正大学)

第 6 回バレーボール学会 (早稲田大学), 2001 年 3 月, p. 17

国際女子バレーボール試合のラインアップ分析に関する研究—'00 オリンピック最終予選, 日本対クロアチア戦の分析—

島津大宣*¹, 泉川喬一*², 山本外憲*³, 田原武彦*⁴, 坂井 充*⁵, 明石正和, 原田 智*⁶ (*¹ 日本女子大学, *² 神奈川工科大学, *³ 杏林大学, *⁴ 奈良大学, *⁵ 九州女子短期大学, *⁶ 立正大学)

第 8 回日本運動・スポーツ科学学会 (日本体育大学健志台), 2001 年 6 月, p. 14

()

-

城西大学理学部研究報告
第 10 卷 (平成 14 年)

編集・発行 城西大学理学部
〒350-0295 埼玉県坂戸市けやき台 1-1
0492-71-7728

編集委員 上原 博通 (委員長)
石橋 宏行 岡田 法雄
宮前 博 明石 正和
加賀美英雄

印刷所 (株) 外為印刷
〒111-0032 東京都台東区浅草 2-29-6
03-3844-3855 (代)
