

Anticancer Res., **19**, 1125-1129 (1999).

Relationship between Radical Intensity and Biological Activity of Cacao Husk Extracts

Noboru Motohashi,^{*1} Masami Kawase (河瀬 雅美),² Teruo Kurihara,³ Yoshiaki Shirataki (白瀧 義明),² Kunio Kamata,⁴ Hideki Nakashima,⁵ Mariappan Premanathan,⁵ Rieko Arakaki,⁵ Kenji Kanbara,⁵ Kazue Satoh,⁶ Hiroshi Sakagami,⁷ Setsuo Saito (齋藤 節生),² and Tetsuo Nakamura⁸

Abstract: The relationship between radical intensity and biological activity of cacao husk extracts was investigated. Electron spin resonance (ESR) spectroscopy demonstrated that the radical intensity of hexane, acetone, methanol and 70% methanol extracts increased with water-solubility. Several fractions of these husk extracts, separated by different column chromatographies, significantly inhibited the cytopathic effect of human immunodeficiency virus (HIV) infection in parallel with their radical intensity. However, their cytotoxic activity against human leukemic and carcinoma cell lines is not always correlated with their radical intensity. Water-soluble and lipophilic compounds might induce cytotoxic activity by different mechanisms.

¹ Meiji Pharmaceutical University, Kiyose-shi, Tokyo

² Faculty of Pharmaceutical Sciences, Josai University, Saitama

³ Faculty of Science, Josai University, Saitama

⁴ Metropolitan Research Laboratory of Public Health, Tokyo

⁵ Department of Microbiology and Immunology, Kagoshima University Dental School, Kagoshima

⁶ Analysis Center, School of Pharmaceutical Sciences, Showa University, Tokyo

⁷ Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama

⁸ Functional Food R and D Laboratories, Meiji Seika Kaisha, Ltd, Saitama, Japan