

Archives of Pharmacal Research, **Vol 22, No 4**, 335-339(1999).

## The Relationship Between Tyrosinase Inhibitory Action and Oxidation-Reduction Potential of Cosmetic Whitening Ingredients and Phenol Derivatives

Katsuya Sakuma, Masayuki Ogawa, Kenji Sugibayashi (杉林 堅次),  
Koh-ichi Yamada (山田 紘一) and Katsumi Yamamoto (山本 勝美)

The oxidation-reduction potentials of cosmetic raw materials, showing tyrosinase inhibitory action, and phenolic compounds structurally similar to L-tyrosine were determined by cyclic voltammetry. The voltammograms obtained could be classified into 4 patterns (patterns 1-4). Pattern 1, characterized by oxidation and reduction peaks as a pair, was observed with catechol, hydroquinone or phenol, and pattern 2 exhibiting another oxidation peak in addition to oxidation and reduction peaks as a pair was found with arbutin, kojic acid, resorcinol, methyl p-hydroxybenzoate and L-tyrosine as the substrate of tyrosinase. Pattern 3 with an independent oxidation peak only was expressed by L-ascorbic acid, and pattern 4 with a reduction peak only at high potentials, by hinokitiol. The tyrosinase inhibitory activity of these compounds was also evaluated using the 50% inhibitory concentration ( $IC_{50}$ ) and the inhibition constant ( $K_i$ ) as parameters. Hinokitiol, classified as pattern 4, showing the highest inhibitory activity (lowest  $IC_{50}$  and  $K_i$ ). Hydroquinone showing the second highest activity belonged to pattern 1, which included compounds showing no inhibition of tyrosinase activity. The inhibitory activity of compounds exhibiting pattern<sup>2</sup> was relatively low with  $K_i$  value being in the order of  $10^{-4}$ M. Although there was no consistent relationship between oxidation-reduction potentials and tyrosinase inhibitory action, the voltammetry data can be used as an additional index to establish the relationship between the structure and the tyrosine inhibitory activity.