

Brain Res., 860, 21-28 (2000).

Relationship between large conductance calcium-activated potassium channel and bursting activity

**Wen Jin (金 文), Aiko Sugaya (菅谷愛子), Tadashi Tsuda (津田 整),
Hiromi Ohguchi (大口広美), and Eiichi Sugaya**

¹Faculty of Pharmaceutical Sciences, Josai University, Keyakidai, Sakado-shi, Saitama-ken, Japan; ²Div., of Oriental Medicine, Tokyo Hospital, School of Medicine, Tokai University, 1-2-5, Yoyogi, Shibuya-ku, Tokyo, 151-0053, Japan

To elucidate the role of the large conductance calcium-activated potassium channel (BK_{Ca} channel) in the production of bursting activity, which is characteristic of convulsions, effects of iberiotoxin (IbTX), a selective blocker of the BK_{Ca} channel, on bursting activity, induced by various procedures were examined using primary cultured neurons from the cerebral cortex of mice. IbTX completely inhibited bursting activity induced by pentylenetetrazol (PTZ), caffeine, inositol 1,4,5,-triphosphate (IP3) and direct forced increase of intracellular calcium. Inherent spontaneous bursting activity in the cerebral cortical neurons of the El mouse, which shows a high susceptibility to convulsions was also completely inhibited by IbTX. Apamin, a specific blocker of the small conductance calcium-activated potassium channel (SK_{Ca} channel) showed no inhibition of bursting activity. These findings suggest that the BK_{Ca} channel is essential for the production of bursting activity, and also suggest the possibility of clinical use of blocking agents of the BK_{Ca} channel against intractable epilepsy.