

Phytotherapy Res., **15**, 337-343 (2001).

Biological activity of Kiwifruit peel extracts

Noboru Motohashi*¹, Yoshiaki Shirataki(白瀧義明)², Masami Kawase(河瀬雅美)², Satoru Tani(谷 覺)², Hiroshi Sakagami³, Kazue Satoh⁴, Teruo Kurihara⁵, Hideki Nakashima⁶, Kristina Wolfard⁷, Csilla Miskolci⁷, and Joseph Molnar⁷

¹Meiji Pharmaceutical University, Tokyo, Japan; ²Faculty of Pharmaceutical Sciences, Josai University, Saitama, Japan; ³Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama, Japan; ⁴Analysis Center, School of Pharmaceutical Sciences, Showa University, Tokyo, Japan; ⁵Faculty of Science, Josai University, Saitama, Japan; ⁶Department of Microbiology and Immunology, Kagoshima University Dental School, Kagoshima, Japan; ⁷Faculty of Medicine, Institute of Microbiology, Albert Szent-Gyorgyi Medical University, Szeged, Hungary

Various bioactive substances in kiwifruit extracts were fractionated by organic solvent extractions, followed by silica gel and ODS chromatographies. Both cytotoxic activity and multi-drug resistance reversal activity were found in the less polar fractions. Cytotoxic activity was not always parallel the radical intensity. Antibacterial activity was distributed into various fractions and all fractions were inactive against *Candida albicans* and *H. pylori*. Only 70% methanol extracts showed anti-human immunodeficiency virus activity, and produced a broad ESR signal under alkaline conditions, in a fashion similar to lignin. These fractions also effectively scavenged O₂⁻ produced by the xanthine-xanthine oxidase reaction, suggesting a bimodal (pro-oxidant and antioxidant) action. These data suggest a medicinal efficacy of kiwifruit peel extracts.