

Chem. Pharm. Bull., 49, 858-862 (2001).

Stilbenoids in Lianas of *Gnetum parvifolium*

Toshiyuki Tanaka¹, Ibrahim Iliya¹, Tetsuro Ito¹, Miyuki Furusawa¹, Ken-ichi Nakaya¹, Munekazu Iinuma¹, Yoshiaki Shirataki(白瀧義明)², Nobuyasu Matsuura³, Makoto Ubukata³, Jin Murata⁴, Fumio Simozono⁴ and Kazunori Hirai⁴

¹Gifu Prefectural Institute of Health and Environmental Sciences, 1-1 Naka Fudogaoka, Kakamigahara, Gifu 504-0838, Japan; ²Faculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai Sakado, Saitama 350-0295, Japan; ³Biotechnology Research center, Toyama Prefectural University, 5180 Kosugi-machi, Izumi-gun, Toyama 939-0398, Japan; ⁴Botanical Gardens, Koishikawa, Graduate School of Science, The University of Tokyo, 3-7-1 Hakusan, Bunkyo-ku, Tokyo 112-0001, Japan

Five new stilbene dimers were isolated from the lianas of *Gnetum parvifolium* in addition to known stilbenoids. The structures of the compounds were established on the basis of spectroscopic evidence, including long-range coupling and nuclear Overhauser effect experiments, in NMR spectrum. Among the isolates, 2b-hydroxyampelopsin F showed potent inhibitory activity in the Maillard reaction.