

Heterocycles, 55, 2123-2130 (2001).

Four New Glucosides of Stilbene Oligomers from the Stem of *Gnetum gnemonoides*

Ibrahim Iliya¹, Toshiyuki Tanaka², Miyuki Furusawa², Zulfiqar Ali², Ken-ichi Nakaya², Munekazu Iinuma², Yoshiaki Shirataki(白瀧義明)³, Jin Murata⁴ and Dedy Darnaedi⁵

¹Gifu Pharmaceutical University, 5-6-1, Mitahora-higashi, Gifu 502-8585, Japan; ²Gifu Prefectural Institute of Health and Environmental Sciences, 1-1 Naka Fudogaoka, Kakamigahara, Gifu 504-0838, Japan; ³Faculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai Sakado, Saitama 350-0295, Japan; ⁴Botanical Gardens, Koishikawa, Graduate School of Science, The University of Tokyo, 3-7-1 Hakusan, Bunkyo-ku, Tokyo 112-0001, Japan; ⁵Indonesian Institute of Sciences, Jalan Ir. H. Juanda 13, Bogor 16122, Indonesia

Four new stilbene oligomeric glucosides, gnemonoside A, B, C and D, were isolated from the stem of *Gnetum gnemonoides* (BRONGN) along with four known stilbenoides, *trans*-resveratrol, gnetins C, D and E. The structures were elucidated on the basis of spectral evidence.