

in vivo, 15, 519-524 (2001).

### Antimicrobial Activity of Prenylflavanones

Sujata G. Dastidar<sup>1</sup>, S. K. Mahapatra<sup>1</sup>, Kumkum Ganguly<sup>1</sup>, A. N. Chakrabarty<sup>2</sup>, Yoshiaki Shirataki(白瀧義明)<sup>3</sup>, and Noboru Motohashi<sup>4</sup>

<sup>1</sup>Division of Microbiology, Department of Pharmaceutical Technology, Jadavpur University, Calcutta 700 032, India; <sup>2</sup>Department of Medical Microbiology and Parasitology, Calcutta University College of Medicine, Calcutta 700 020, India; <sup>3</sup>Faculty of Pharmaceutical Sciences, Josai University, Sakado, Saitama 350-0295, Japan; <sup>4</sup>Meiji Pharmaceutical University, Kiyose, Tokyo 204-8588, Japan

Among two flavanones [YS01, YS02] and eight prenylflavanones [YS03-YS10], preliminary screening with fifteen test bacterial strains showed that YS06 was the most active agent. YS06 exhibited highly significant antimicrobial action when tested against 228 bacterial strains comprising two Gram-positive and six Gram-negative genera. The in vitro susceptibility test was carried out by determining the minimum inhibitory concentration (MIC) of YS06 by agar dilution technique. Twenty-two out of fifty strains of *Staphylococcus aureus* were inhibited at 25 to 50  $\mu$ g/mL of the agent. YS06 also inhibited strains of *Salmonella*, *Shigella* and a few strains of *Escherichia coli* strains were also highly sensitive to YS06, while *Klebsiella* spp. and *Pseudomonas aeruginosa* were much less sensitive. In *in vivo* study, YS06 offered significant protection ( $p < 0.001$  according to chi-square test) to Swiss albino mice (challenged with 50 minimum lethal dose (MLD, virulent bacterium) at concentrations of 160 and 80  $\mu$ g/mouse.