

J.Org.Chem., 66, 7107-7112 (2001).

Total Synthesis of (\pm)-Stemodinone via an Efficient Ring-Exchange Strategy

Tetsuaki Tanaka¹, Kazuo Murakami¹, Atsushi Kanda¹, Debasis Patra¹, Sachiko Yamamoto¹, Norifumi Satoh¹, Sang-Won Kim(金相元)², S. M. Abdur Rahman¹, Hiroaki Ohno¹, and Chuzo Iwata¹

¹Graduate school of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan, and ²Faculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai, Sakado, Saitama 354-0295, Japan

A total synthesis of (\pm)-Stemodinone, a tetracyclic stemodane diterpene, from the known tricyclic methyl olefin 11 is described. The key steps involve an efficient ring-exchange reaction and palladium(0)-catalyzed lactone migration. The ring-exchange strategy for controlling the stereochemistry was based on an initial Diels-Alder reaction to form a new ring followed by cleavage of the original ring. Cleavage of the original ring of the Diels-Alder adduct 9 was achieved by an initial regio- and chemoselective Baeyer-Villiger oxidation followed by the Pd(0)-catalyzed lactone_migration reaction reported by us.

