

Chem. Pharm. Bull., 50, 796-801 (2002).

Stilbene Derivatives from Two Species of Gnetaceae

Ibrahim Iliya^a, Toshiyuki Tanaka^b, Munekazu Iinuma^b, Zulfiqur Ali^b, Miyuki Furusawa^b, Ken-ichi Nakaya^b, Yoshiaki Shirataki (白瀧 義明)^b, Jin Murata^b and Dedy Darnaed^b

^aGifu Pharmaceutical University, 5-6-1 Mitahora-higashi, Gifu 502-8585, Japan;

^bGifu Prefectural Institute of Health and Environmental Sciences, 1-1 Naka Fudogaoka, Kakamigahara, Gifu 504-0838, Japan; ^cFaculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai Sakado, Saitama 350-0295, Japan;

^dBotanical Gardens, Koishikawa, Graduate School of Science, The University of Tokyo, 3-7-1 Hakusan, Bunkyo-ku, Tokyo 112-0001, Japan; ^eIndonesian Institute of Sciences, Jalan Ir. H. Juanda 13, Bogor 16122, Indonesia

Five new stilbene oligomers (gnemonols A, B and C, gnemonoside E and gnatal) were isolated together with 2b-hydroxyampelopsin F and gnetin E from *Gnetum gnemon* and *G. gnemonoides*. The structures of the compounds were elucidated on the basis of spectral evidence.