

Oriental Pharm. Experimental Med., 3, 100-105 (2003).

**Multidrug resistance reversal in mouse lymphoma cells by Indian tea leaves, Indian coffee seeds and chicory.**

Rao BK, <sup>1</sup> Motohashi N, <sup>2</sup> Kawase M (河瀬雅美), <sup>3</sup> Spengler G, <sup>4</sup> Molnar J <sup>4</sup>

<sup>1</sup>Department of Chemistry, Nagarjuna University, Andhra Pradesh, India; <sup>2</sup>Meiji Pharmaceutical University, Tokyo, Japan; <sup>3</sup>Faculty of Pharmaceutical Sciences, Josai University, Saitama, Japan; <sup>4</sup>Faculty of Medicine, Albert Szent-Gyorgyi Medical University, Szeged, Hungary

Systematic analysis of caffeine from the commercial samples of India tea leaves was performed by a routine method and the content of caffeine was found to be 19.0-37.4 mg/100 g leaves. The caffeine contents from coffee seeds and chicory from Indian origin were analyzed and found to be 0.6540-1.4920 g/100 g seeds. Caffeine contents of roasted Indian chicory roots were lower than either those of Indian tea leaves or Indian coffee seeds. The multidrug resistance (MDR) reversing effects were tested on a mouse leukemia cell line of L-5178 cells by methanol extracts [M1-M15] of Indian tea leaves and coffee seeds, comparing to a control of verapamil. The effects were measured by fluorescence ratio between treated and untreated group cells. Among fifteen methanol extracts, a Gemini tea [M6](fluorescence activity ratio 5.26) had the most potent effect for L-5178 cells. The extract M6 was 0.63-fold of verapamil. We suggest that one of mechanisms of reversal by M6 might have strong affinity to dopamine D<sub>1</sub> and D<sub>2</sub> receptors. Further studies with many more tumor and normal cell lines are necessary to confirm the MDR reversal specificity of coffee methanol extracts.