

Chem. Pharm. Bull., 51, 85-88 (2003).

**Three New Trimeric Stilbenes from *Gnetum gnemon*.**

Ibrahim Iliya<sup>1</sup>, Zulfiqar Ali<sup>2</sup>, Toshiyuki Tanaka<sup>2</sup>, Munekazu Iinuma<sup>1</sup>, Miyuki Furasawa<sup>2</sup>, Ken-ichi Nakaya<sup>2</sup>, Yoshiaki Shirataki(白瀧義明)<sup>3</sup>, Jin Murata<sup>4</sup>, Dedy Darnaedi<sup>5</sup>, Nobuyasu Matsuura<sup>6</sup>, and Makoto Ubukata<sup>6</sup>

<sup>1</sup>Gifu Pharmaceutical University, 5-6-1 Mitahora-higashi, Gifu 502-8585, Japan, <sup>2</sup>Gifu Prefectural Institute of Health and Environmental Sciences, 1-1 Naka-fudogaoka, Kagamigahara 504-0838, Japan, <sup>3</sup>Faculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai, Sakado, Saitama 350-0295, Japan, <sup>4</sup>Botanical Gardens, Koishikawa, Graduate School of Science, University of Tokyo, 3-7-1 Hakusan, Bunkyo-ku, Tokyo 112-0001, Japan, <sup>5</sup>Indonesian Institute of Sciences, Jalan Ir. H. Juanda 13, Bogor 16122, Indonesia, <sup>6</sup>Toyama Prefectural University, 5180 Kosugi-machi, Izumi-gun, Toyama 939-0398, Japan

Three stilbene trimers (gnemonols D, E, F) were isolated from the root of *Gnetum gnemon*. The structures were determined by spectroscopic analysis. In addition, the antioxidant activity of the compounds on lipid peroxide inhibition and super oxide scavenging activity were also investigated.