

Tetrahedron, 59, 5347-5393 (2003).

New resveratrol oligomers in the stem bark of *Vatica pauciflora*

Tetsuro Ito¹, Toshiyuki Tanaka¹, Munekazu Iinuma², Ibrahim Iliya², Ken-ichi Nakaya¹, Zulfiqar Ali¹, Yoshiakazu Takahashi³, Ryuichi Sawa³, Yoshiaki Shirataki (白瀧義明)⁴, Jin Murata⁵ and Dedy Darnaedi⁶

¹Gifu Prefectural Institute of Health and Environmental Sciences, Naka-fudogaoka, Kakamigahara, Gifu 504-0838, Japan, ²Gifu Pharmaceutical University, 5-6-1 Mitahora-higashi, Gifu 502-8585, Japan, ³Institute of Microbial Chemistry, 3-14-23 Kamiosaki, Shinagawa-ku, Tokyo 141-0021, Japan, ⁴Faculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai Sakado, Saitama 350-0295, Japan, ⁵Botanical Gardens, Koishikawa, Graduate School of science, University of Tokyo, 3-7-1, Hakusan, Bunkyo-Ku, Tokyo, 112-0001, Japan, ⁶Indonesian Institute of Sciences, Jalan Ir. H.Juanda 13, Bogor 16122, Indonesia

Five new resveratrol oligomers; pauciflorols A-C (1-3), Isovaticanols B (6) and C (8), and three new oligostibene glucosides; pauciflorosides A (11), B(13), C(14), were isolated from the stem bark of *Vatica pauciflora* (Dipterocarpaceae) together with known 17 resveratrol oligomers (4, 5, 7, 9, 10, 12 and 15-25) and bergenin (26). The structures of isolates were established in the basis of detailed spectroscopic analysis. The typical and characteristic spectral properties of some resveratrol oligomers were also discussed.